

# Transgenerational effects of global environmental change: long-term CO<sub>2</sub> and nitrogen treatments influence offspring growth response to elevated CO<sub>2</sub>

Jennifer A. Lau · Jill Peiffer ·  
Peter B. Reich · Peter Tiffin

Received: 17 January 2008 / Accepted: 29 July 2008 / Published online: 21 August 2008  
© Springer-Verlag 2008

**Abstract** Global environmental changes can have immediate impacts on plant growth, physiology, and phenology. Long-term effects that are only observable after one or more generations are also likely to occur. These transgenerational effects can result either from maternal environmental effects or from evolutionary responses to novel selection pressures and are important because they may alter the ultimate ecological impact of the environmental change. Here, we show that transgenerational effects of atmospheric carbon dioxide (CO<sub>2</sub>) and soil nitrogen (N) treatments influence the magnitude of plant growth responses to elevated CO<sub>2</sub> (eCO<sub>2</sub>). We collected seeds from *Lupinus perennis*, *Poa pratensis*, and *Schizachyrium scoparium* populations that had experienced five growing seasons of ambient CO<sub>2</sub> (aCO<sub>2</sub>) or eCO<sub>2</sub> treatments and ambient or increased N deposition and planted these seeds into aCO<sub>2</sub> or eCO<sub>2</sub> environments. We found that the offspring eCO<sub>2</sub> treatments stimulated immediate increases in *L. perennis* and *P. pratensis* growth and that the maternal CO<sub>2</sub> environment influenced the magnitude of this growth response for

*L. perennis*: biomass responses of offspring from the eCO<sub>2</sub> maternal treatments were only 54% that of the offspring from the aCO<sub>2</sub> maternal treatments. Similar trends were observed for *P. pratensis* and *S. scoparium*. We detected some evidence that long-term N treatments also altered growth responses to eCO<sub>2</sub>; offspring reared from seed from maternal N-addition treatments tended to show greater positive growth responses to eCO<sub>2</sub> than offspring from ambient N maternal treatments. However, the effects of long-term N treatments on offspring survival showed the opposite pattern. Combined, our results suggest that transgenerational effects of eCO<sub>2</sub> and N-addition may influence the growth stimulation effects of eCO<sub>2</sub>, potentially altering the long-term impacts of eCO<sub>2</sub> on plant populations.

**Keywords** Climate change · Evolution · FACE · Maternal effect

## Introduction

The environment is rapidly changing on a global scale, largely as a result of anthropogenic pressures. Plant populations are likely to respond to global changes in the short-term with plastic changes in phenotype. For example, numerous studies have documented immediate increases in plant growth in response to elevated atmospheric carbon dioxide (eCO<sub>2</sub>) concentrations and nitrogen (N) fertilization (Bazzaz 1990; Reich et al. 2001a; Poorter and Navas 2003; Niklaus and Körner 2004; Reich et al. 2006). Long-term responses that only become apparent after one or more generations are also possible. These transgenerational effects can result from maternal environmental effects, which occur when the environmental conditions experienced by parents influence offspring traits (Roach and

---

Communicated by Manuel Lerdau.

---

J. A. Lau · J. Peiffer · P. Tiffin  
Department of Plant Biology, University of Minnesota,  
St Paul, MN 55108, USA

P. B. Reich  
Department of Forest Resources, University of Minnesota,  
St Paul, MN 55108, USA

*Present Address:*  
J. A. Lau (✉)  
Kellogg Biological Station and Department of Plant Biology,  
Michigan State University, 3700 East Gull Lake Dr.,  
Hickory Corners, MI 49060, USA  
e-mail: jenlau@msu.edu

Wulff 1987; Rossiter 1996), or from genetic changes in response to altered patterns of natural selection. Regardless of the underlying cause, transgenerational effects may either increase (e.g., Bezemer et al. 1998; Reale et al. 2003) or decrease (e.g., Huxman et al. 1998, 2001) the initial phenotypic response to the environmental change. Thus, the two types of transgenerational effects (maternal environmental and genetic) may alter the long-term predicted effects of environmental change on natural communities.

Maternal environmental effects commonly occur in both plant and animal species when the maternal environment influences resource allocation to offspring. One example of this is when parents occupy resource-rich habitats: offspring size and subsequent growth and fitness are often increased relative to the offspring of parents inhabiting more stressful environments (e.g., Durrant 1958; Aarssen and Burton 1990; Trombulak 1991; reviewed in Rossiter 1996). Maternal environmental conditions can also influence a wide range of other offspring characters, including anti-predator defenses (Agrawal et al. 1999) and physiological traits (Huxman et al. 2001; reviewed in Rossiter 1996). As a result, maternal effects can impact population dynamics (e.g., Ginzburg and Taneyhill 1994), responses to abiotic stress (Aarssen and Burton 1990), and interactions with other community members, such as herbivores and competitors (Agrawal et al. 1999; Platenkamp and Shaw 1993). Maternal environmental effects also can influence plastic responses to variation in the offspring environment (e.g., Aarssen and Burton 1990; Bezemer et al. 1998; Huxman et al. 2001). For example, offspring from maternal plants that had experienced nutrient stress were smaller and germinated later in both low and high nutrient environments, but they also exhibited greater tolerance to low nutrient conditions (Aarssen and Burton 1990). In another example, offspring of plants grown in eCO<sub>2</sub> environments showed an increased growth response to eCO<sub>2</sub> compared to offspring from plants grown in ambient CO<sub>2</sub> (aCO<sub>2</sub>) environments (Bezemer et al. 1998). As these studies illustrate, the maternal effect can result in reduced *or* heightened plastic responses to offspring environmental conditions.

Populations also might evolve in response to the novel selection pressures that may accompany environmental change, resulting in a very different type of long-term effect. While maternal environmental effects typically decrease over time and often become non-detectable after one or a few generations or even over the course of the offspring life cycle (Roach and Wulff 1987), genetic changes in response to novel environmental conditions are likely to persist longer, even if organisms are no longer exposed to that environment. Adaptive evolutionary changes are expected to reduce any negative fitness effects of the environmental perturbation; however, these evolutionary changes also have the potential to influence population dynamics (e.g., Yoshida et al. 2003,

2004), ecosystem processes (Wieneke et al. 2004; Collins et al. 2006), interactions with other community members (Snaydon and Davies 1982; Lau et al. 2008), and, most importantly, the magnitude of response to the environmental change itself (Snaydon 1970; Snaydon and Davies 1972, 1982) (reviewed in Hairston et al. 2005).

We have assessed how transgenerational effects of long-term CO<sub>2</sub> and N treatments influence the magnitude of the growth response to eCO<sub>2</sub> concentrations. We first tested how offspring eCO<sub>2</sub> treatments impact plant growth and survival and then examined how maternal CO<sub>2</sub> and N environments affect plant growth and growth responses to offspring CO<sub>2</sub> treatments. Finally, we investigated how maternal CO<sub>2</sub> and N treatments impact resource allocation to seeds and whether differences in seed mass can explain any of the observed transgenerational effects.

## Materials and methods

### Study system and experimental design

We investigated transgenerational effects of eCO<sub>2</sub> and soil N deposition on the growth of *Schizachyrium scoparium* (Poaceae), *Poa pratensis* (Poaceae), and *Lupinus perennis* (Fabaceae). These three species represent different functional groups (C<sub>4</sub> grass, C<sub>3</sub> grass, and legume, respectively) that are expected to respond differently to eCO<sub>2</sub> and N-addition. We performed a partial reciprocal transplant experiment in which we planted seeds collected from populations growing in long-term atmospheric CO<sub>2</sub> and soil N treatments into aCO<sub>2</sub> and eCO<sub>2</sub> environments. Long-term CO<sub>2</sub> and N treatments were manipulated in a modified split-plot design as part of the BioCON experiment at Cedar Creek Natural History Area, Minnesota, USA (Reich et al. 2001a). In the BioCON experiment, biodiversity (plant species richness), CO<sub>2</sub>, and N are all experimentally manipulated. The two CO<sub>2</sub> treatments (aCO<sub>2</sub> approx. 370 μmol/mol; eCO<sub>2</sub> approx. 560 μmol/mol) were applied to whole plots (three 20-m-diameter rings per treatment) using Free Air CO<sub>2</sub> Enrichment. Nitrogen treatments (0 or 4 g N m<sup>2</sup>/year) were applied to half of the 2 × 2-m sub-plots within each whole plot (i.e., ring). Carbon dioxide and N treatments have been applied annually since 1998. Only seeds from the high-diversity (16 plant species) treatment plots were used in this study ( $n = 48$  plots, four plots per N treatment per ring). In 1997, each high-diversity plot was seeded with 12 g/m<sup>2</sup> of seed, divided equally among 16 species that were native or naturalized to the site. All seeds were bulked collections, obtained from local nurseries: *S. scoparium* seed were obtained from Prairie Restorations (Princeton, MN); *L. perennis* seeds were obtained from the Prairie Moon Nursery (Winona, MN).

Seeds used in this study were collected from a single maternal plant per species from each plot, resulting in 12 half-sib families from each maternal CO<sub>2</sub> × N treatment. All seeds were collected in the fall of 2002, after treatments had been applied for five full growing seasons. Even though all plants used in this experiment are perennial species, because plots were initially established from seed, evolutionary responses could result from differential survival of the initial stand of seedlings as well as from genotypic differences in sexual reproduction and recruitment from seed in subsequent years.

#### Planting design and plant measurements

Seeds were initially planted into 164-ml conetainers (Ray Leach Conetainers; Stuewe & Sons, Corvallis, OR) that had been filled with potting mix (Sunshine Mix #5; Sun Gro Horticulture Canada, Spruce Grove, AB) and bottom-watered until saturated. For *S. scoparium* and *L. perennis*, we planted four seeds (all from a single maternal family) per conetainer on 24 April 2006, placed the conetainers in a dark cold-room for 8 days at 4°C, and then moved them to a greenhouse (ambient CO<sub>2</sub>) where they remained until the seeds began to germinate. For *L. perennis*, we planted one seed per conetainer on 5 May 2006. *Lupinus perennis* seeds were scarified and soaked overnight prior to planting to improve germination, but *L. perennis* individual plants were not inoculated with rhizobia. Because we germinated all plants under aCO<sub>2</sub> conditions, we unfortunately missed eCO<sub>2</sub> effects on germination and very early seedling growth; however, germinating the plants in this way was necessary for facilitating the germination of a fourth species used in a separate experiment conducted simultaneously. All plants were moved to the field and placed into the appropriate offspring CO<sub>2</sub> treatment in the BioCON facility on 7 May 2006. We placed two conetainers from each maternal family into each ring (six conetainers per family per offspring CO<sub>2</sub> treatment). A total of 1728 conetainers were planted (12 conetainers per family × 48 maternal families per species × 3 species); however, because of poor germination, which was likely due to the poor quality and limited numbers of some seed, actual sample sizes were much lower (final *n* = 236, 135, or 264 conetainers for *L. perennis*, *S. scoparium*, and *P. pratensis*, respectively). Sample sizes were roughly equivalent across most maternal environment × offspring environment treatment combinations for most species, and there were no missing cells (i.e., treatment combinations with no data; *n* = 20–41 plants/treatment combination for *L. perennis*; *n* = 25–39 for *P. pratensis*, and *n* = 7–26 for *S. scoparium*). However, we report statistics based on Type 3 sums of squares to further minimize biases due to unequal sample sizes between treatments (Shaw and Mitchell-Olds 1993).

We weighed each individual *L. perennis* seed prior to planting. Additionally, we obtained family mean seed mass for *S. scoparium* and *P. pratensis* by weighing all collected seeds from a given maternal plant and dividing by seed number. For all three species, we recorded emergence date, and on 10 July 2006 we measured the height of the tallest culm and culm number for *S. scoparium* and *P. pratensis* and leaf number and leaf size (length of longest leaflet) for *L. perennis*. After 3 months of growth, when foliage was beginning to turn brown, we harvested aboveground biomass. For *S. scoparium* and *L. perennis*, we also harvested the belowground biomass (*P. pratensis* roots were difficult to separate from the soil) and measured the length of longest root (*L. perennis*) or belowground biomass (*S. scoparium*). All biomass was dried at 60°C for at least 2 days prior to weighing. Because four grass seeds were planted into each conetainer, many conetainers contained multiple individuals; all plants in a conetainer were weighed together to obtain total biomass, and heights and culm numbers were averaged for all analyses. Survival was assessed at the final harvest.

#### Statistical analysis

All analyses were performed using SAS 2001 statistical software (SAS Institute, Cary N.C.).

We used mixed model multivariate analysis of variance (MANOVA; PROC GLM) and ANOVA (PROC MIXED) to determine how both the offspring and maternal growing environments influenced plant growth and plant responsiveness to eCO<sub>2</sub>. Data for each species were analyzed separately. The N and CO<sub>2</sub> treatments that the seeds were collected from (“maternal N treatment” and “maternal CO<sub>2</sub> treatment”), the CO<sub>2</sub> treatment in which the experimental plants were reared (“offspring CO<sub>2</sub> treatment”), and all interactions were included in the model as fixed factors. Ring nested within the offspring CO<sub>2</sub> treatment and maternal family nested within the maternal N × maternal CO<sub>2</sub> interaction were included as random factors. Emergence date and seed mass were included as covariates in the *L. perennis* analyses. Mean emergence date, family mean seed mass, and number of individuals per conetainer were included as covariates in the *P. pratensis* and *S. scoparium* analyses. Preliminary analyses revealed that seed mass did not significantly influence any *P. pratensis* or *S. scoparium* growth measures; therefore, it was not included in the final analyses. Results from models including covariates were qualitatively similar to results from additional analyses that did not include the seed character covariates, so only results from the full model are presented. For *L. perennis*, final aboveground biomass, root length, leaf number, and leaf length were included as response variables. For the grasses, final aboveground biomass, root biomass (*S. scoparium*

only), culm number, and height were included as response variables.

The effects of maternal CO<sub>2</sub> and N treatments and offspring CO<sub>2</sub> treatment on survival were tested with logistic ANOVA (PROC GENMOD). In these analyses, maternal CO<sub>2</sub> treatment, maternal N treatment, offspring CO<sub>2</sub> treatment, and all interactions were included as predictor variables, and the proportion of germinants in each pot surviving to harvest was the response variable. For *L. perennis*, however, survival was scored as a binomial response variable since only one seed was planted per pot. Emergence date and seed mass were included as covariates. As above, data for each species were analyzed separately.

Because maternal environmental effects often result from differences in resource allocation to offspring, we used ANOVA (PROC MIXED) to investigate how maternal CO<sub>2</sub> and N treatments influenced the mass of field-collected seeds used in the reciprocal transplant experiment. For *L. perennis*, the model included maternal CO<sub>2</sub> and maternal N treatments as fixed factors, and maternal family nested within the maternal CO<sub>2</sub> × maternal N interaction as a random factor. Individual seed mass was the response variable. For *S. scoparium* and *P. pratensis*, the mean family seed mass was the response variable, and maternal CO<sub>2</sub> and maternal N treatments were included as predictor variables.

## Results

### Direct phenotypic effects of CO<sub>2</sub>

Compared to aCO<sub>2</sub> treatments, offspring eCO<sub>2</sub> treatments increased *L. perennis* biomass by 42%, leaflet length by 17%, and leaf number by 12% (Table 1; Fig. 1a). Similarly, eCO<sub>2</sub> increased *P. pratensis* biomass, height, and culm number by 116, 44, and 31%, respectively (Table 1; Fig. 1b). In contrast, *S. scoparium*, the C<sub>4</sub> grass, did not respond significantly to eCO<sub>2</sub> (Table 1; Fig. 1c). These results are consistent with prior results for these three species in BioCON (Reich et al. 2001b) as well as with those from other studies demonstrating that species in the C<sub>4</sub> functional group tend to show weaker responses to eCO<sub>2</sub> (Poorter and Navas 2003). On average, offspring eCO<sub>2</sub> treatments increased *P. pratensis* survival by 21% but did not significantly influence *L. perennis* or *S. scoparium* survival ( $P > 0.3$ ) (Table 2; Fig. 2).

### Transgenerational effects of CO<sub>2</sub> and N

#### *Effects of maternal CO<sub>2</sub> environment*

Offspring response to eCO<sub>2</sub> was strongly influenced by the maternal CO<sub>2</sub> environment in *L. perennis* (significant

maternal CO<sub>2</sub> × offspring CO<sub>2</sub> interaction; Table 1). Offspring from aCO<sub>2</sub> maternal treatments showed greater biomass increases in response to eCO<sub>2</sub> than offspring from eCO<sub>2</sub> maternal treatments (Fig. 3). While this difference was only statistically significant for *L. perennis*, *P. pratensis* and *S. scoparium* showed similar trends (Fig. 3). The biomass stimulation effect of offspring eCO<sub>2</sub> treatments on plants from eCO<sub>2</sub> maternal environments (averaged across N treatments) was only 54, 26, or 72% that of offspring from aCO<sub>2</sub> maternal treatments for *L. perennis*, *S. scoparium*, and *P. pratensis*, respectively (Fig. 3). A significant maternal CO<sub>2</sub> environment × offspring CO<sub>2</sub> environment interaction also was detected for *P. pratensis* culm number in the univariate analysis, where genotypes from eCO<sub>2</sub> maternal treatments showed only 34% of the growth response to offspring eCO<sub>2</sub> compared to genotypes from aCO<sub>2</sub> maternal treatments (Table 1). The maternal CO<sub>2</sub> environment had minimal impacts on other traits (Table 1). We also did not detect any significant effects of maternal CO<sub>2</sub> environment or interactions between maternal CO<sub>2</sub> and offspring CO<sub>2</sub> treatments on the survival of any species (Table 2).

#### *Effects of maternal N environment*

Although the multivariate analyses revealed little evidence that maternal N environment affects *L. perennis* and *S. scoparium* growth or growth responses to eCO<sub>2</sub>, maternal N environment × offspring CO<sub>2</sub> environment interactions were detected for *L. perennis* root length and for *S. scoparium* height (Table 1). For both species, offspring from maternal N-addition treatments exhibited greater positive growth responses to offspring eCO<sub>2</sub> treatments. *Lupinus perennis* individuals reared from seed collected from ambient N maternal treatments had shorter root lengths when reared under eCO<sub>2</sub> than when reared under aCO<sub>2</sub>, but root lengths of plants reared from N-addition maternal treatments responded positively to eCO<sub>2</sub> offspring treatments (a 3% decrease compared to a 9% increase). Similarly, *S. scoparium* from ambient maternal N environments responded to offspring eCO<sub>2</sub> treatments with decreases in height (a 6% decrease), but individuals from N-addition maternal environments were 16% taller when grown in offspring eCO<sub>2</sub> than when grown in the aCO<sub>2</sub> treatments. Greater positive biomass responses to eCO<sub>2</sub> for *S. scoparium* from N-addition maternal treatments were also observed (Fig. 1c), although this interaction was not statistically significant ( $P > 0.15$ ). Additionally, maternal N and maternal CO<sub>2</sub> treatments interacted to influence the growth response of *P. pratensis* to offspring eCO<sub>2</sub> treatments. A significant three-way interaction between maternal CO<sub>2</sub> environment, maternal N environment, and offspring CO<sub>2</sub> environment was detected in the multivariate analysis of

**Table 1** Effects of maternal CO<sub>2</sub> and N treatments, offspring CO<sub>2</sub> treatment and all interactions on *Lupinus perennis*, *Poa pratensis*, and *Schizachyrium scoparium* growth measures


<i>L. perennis</i>	MANOVA			Biomass	Root length	Leaf length	Leaf number
	Pillai	df	F value	F value	F value	F value	F value
CO <sub>2</sub> Offspring (O)	0.98	4,1	15.99	79.02****	0.89	92.12****	9.60*
CO <sub>2</sub> Maternal (M)	0.13	4,37	1.36	3.33	0.02	0.23	0.24
N <sub>Maternal</sub>	0.11	4,37	1.15	1.10	0.12	0.48	1.05
CO <sub>2</sub> Off × CO <sub>2</sub> M	0.15	4,72	3.26*	7.24**	0.58	0.56	1.32
CO <sub>2</sub> Off × N <sub>M</sub>	0.06	4,72	0.31	0.04	6.55*	2.40	1.64
CO <sub>2</sub> M × N <sub>M</sub>	0.06	4,37	0.58	0.90	2.21	1.87	0.17
CO <sub>2</sub> Off × CO <sub>2</sub> M × N <sub>M</sub>	0.05	4,72	1.04	0.65	0.12	0.19	2.82
Germination date	0.15	4,63	2.78*	40.37****	0.49	6.39*	20.17****
Seed mass	0.13	4,63	2.27	7.86**	0.73	0.01	2.28
Ring(CO <sub>2</sub> Off)	0.19	16,264	0.82	χ <sup>2</sup> = 0.0	χ <sup>2</sup> = 1.0	χ <sup>2</sup> = 0.0	χ <sup>2</sup> = 0.1
Family(CO <sub>2</sub> M × N <sub>M</sub> )	2.17	160,264	1.95****	χ <sup>2</sup> = 4.4*	χ <sup>2</sup> = 2.6	χ <sup>2</sup> = 12.0****	χ <sup>2</sup> = 10.2****
<i>P. pratensis</i>	MANOVA			Biomass	Culm height	Culm number	
	Pillai	df	F value	F value	F value	F value	
CO <sub>2</sub> Offspring (O)	0.95	3,2	11.96	41.36**	41.51****	8.30*	
CO <sub>2</sub> Maternal (M)	0.05	3,35	0.60	0.03	0.03	0.00	
N <sub>Maternal</sub>	0.01	3,35	0.17	0.45	0.38	0.02	
CO <sub>2</sub> Off × CO <sub>2</sub> M	0.08	3,57	1.59	1.51	0.00	4.00*	
CO <sub>2</sub> Off × N <sub>M</sub>	0.01	3,57	0.27	0.04	0.06	1.33	
CO <sub>2</sub> M × N <sub>M</sub>	0.07	3,35	0.94	4.12*	0.12	3.91	
CO <sub>2</sub> Off × CO <sub>2</sub> M × N <sub>M</sub>	0.13	3,57	2.88*	0.89	0.41	0.04	
Germination date	0.19	3,17	1.30	41.37****	3.97*	29.74****	
Plants per pot	0.75	3,17	17.13****	22.76****	1.00	3.45	
Ring(CO <sub>2</sub> Off)	0.71	12,57	1.47	χ <sup>2</sup> = 0.1	χ <sup>2</sup> = 0.0	χ <sup>2</sup> = 1.0	
Family(CO <sub>2</sub> M × N <sub>M</sub> )	2.22	111,57	1.46	χ <sup>2</sup> = 2.2	χ <sup>2</sup> = 4.6*	χ <sup>2</sup> = 0.0	
<i>S. scoparium</i>	MANOVA			Biomass	Root mass	Culm height	Culm number
	Pillai	df	F value	F value	F value	F value	F value
CO <sub>2</sub> Offspring (O)	0.80	4,1	1.00	1.04	0.04	0.79	0.00
CO <sub>2</sub> Maternal (M)	0.15	4,30	1.34	0.00	0.14	0.00	1.09
N <sub>Maternal</sub>	0.06	4,30	0.47	2.76	0.37	1.61	0.25
CO <sub>2</sub> Off × CO <sub>2</sub> M	0.06	4,32	0.55	0.48	0.37	0.72	0.63
CO <sub>2</sub> Off × N <sub>M</sub>	0.05	4,32	0.42	2.09	0.01	4.36*	1.01
CO <sub>2</sub> M × N <sub>M</sub>	0.04	4,30	0.19	1.15	0.55	0.35	0.02
CO <sub>2</sub> Off × CO <sub>2</sub> M × N <sub>M</sub>	0.08	4,32	0.67	0.00	0.15	0.11	0.25
Germination date	0.65	4,11	5.02*	0.34	0.81	0.16	0.09
Plants per pot	0.65	4,11	5.18*	14.21***	10.24**	0.02	7.16**
Ring(CO <sub>2</sub> Off)	1.19	16,56	1.49	χ <sup>2</sup> = 0.2	χ <sup>2</sup> = 0.3	χ <sup>2</sup> = 0.0	χ <sup>2</sup> = 0.0
Family(CO <sub>2</sub> M × N <sub>M</sub> )	3.02	132,56	1.30	χ <sup>2</sup> = 0.1	χ <sup>2</sup> = 0.0	χ <sup>2</sup> = 3.0*	χ <sup>2</sup> = 0.0

MANOVA, Multivariate analysis of variance

\*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$ , \*\*\*\*  $P < 0.0001$

*P. pratensis* growth traits (Table 1), but because no significant three-way interactions were detected in the univariate analyses, this effect was likely due to joint effects on multiple traits. In general, the effects of maternal CO<sub>2</sub> environment

on the response to offspring CO<sub>2</sub> environment tended to be greater for *P. pratensis* from N-addition maternal environments. For *P. pratensis* from ambient N maternal treatments, plants from aCO<sub>2</sub> and eCO<sub>2</sub> maternal treatments



**Fig. 1** Effects of offspring CO<sub>2</sub> treatments on aboveground biomass of *Lupinus perennis* (a), *Poa pratensis* (b), or *Schizachyrium scoparium* (c) populations collected from maternal aCO<sub>2</sub> (open symbols) vs. eCO<sub>2</sub> (filled symbols) and N-addition (squares) vs. ambient N (circles) treatments. Values shown are least-squares means  $\pm$  1SE. Symbols are offset so that non-overlapping error bars can be easily observed

showed similar increases in biomass in response to offspring eCO<sub>2</sub> treatments. For *P. pratensis* from N-addition maternal treatments, however, the growth stimulation effects of offspring eCO<sub>2</sub> treatments were greater on plants from aCO<sub>2</sub> maternal treatments than on those from eCO<sub>2</sub> maternal treatments, and a marginally significant maternal CO<sub>2</sub>  $\times$  offspring CO<sub>2</sub> interaction on biomass was detected for offspring from the maternal N-addition treatment ( $F_{1,54} = 3.55$ ,  $P = 0.065$ ), but not for offspring from the ambient N maternal treatment ( $F_{1,60} = 0.08$ ,  $P = 0.78$ ).

Averaged across offspring and maternal CO<sub>2</sub> treatments, *L. perennis* offspring reared from seeds collected from N-addition maternal environments had a lower survival than offspring from seeds from ambient N maternal environments (mean  $\pm$  SD: N-addition  $0.81 \pm 0.40$ ; ambient N  $0.90 \pm 0.30$ , Table 2). *Poa pratensis* survival showed similar, but non-significant, patterns (N-addition  $0.48 \pm 0.49$ ; ambient N  $0.60 \pm 0.49$ ,  $P < 0.12$ ). No significant main effect of maternal N environment on survival was detected for *S. scoparium* ( $P > 0.5$ ); however, we detected significant interactions between maternal N environment and offspring CO<sub>2</sub> environment on *S. scoparium* survival (Table 2). Genotypes from maternal N-addition treatments showed reduced survival in eCO<sub>2</sub> offspring environments compared to aCO<sub>2</sub> offspring environments, whereas genotypes from ambient N maternal treatments showed increased survival in offspring eCO<sub>2</sub> treatments (Fig. 2c). Furthermore, these effects were heightened when plants were collected from eCO<sub>2</sub> maternal environments (Fig. 2c; significant CO<sub>2</sub> offspring  $\times$  CO<sub>2</sub> maternal  $\times$  N maternal interaction, Table 2).

#### Effects of maternal CO<sub>2</sub> and N environments on seed mass

We did not detect any evidence that maternal N treatment affected seed mass of our study species; however, maternal CO<sub>2</sub> treatment affected *L. perennis* and *S. scoparium* seed mass. *Lupinus perennis* seeds from eCO<sub>2</sub> maternal environments were heavier than those from aCO<sub>2</sub> maternal environments (mean seed mass  $\pm$  SE: aCO<sub>2</sub>  $8.0 \pm 0.9$  mg; eCO<sub>2</sub>  $11.0 \pm 0.9$  mg;  $F_{1,44} = 6.13$ ,  $P = 0.017$ ). *Schizachyrium scoparium* showed the opposite pattern, and seeds from aCO<sub>2</sub> environments were heavier than seeds from eCO<sub>2</sub> environments (mean seed mass  $\pm$  SE: aCO<sub>2</sub>  $1.6 \pm 0.06$  mg; eCO<sub>2</sub>  $1.3 \pm 0.07$  mg;  $F_{1,5} = 7.62$ ,  $P = 0.04$ ). Differences in seed mass that resulted from the maternal environment did not, however, explain the differences in offspring responses to CO<sub>2</sub>. For *L. perennis*, seed mass was positively correlated with biomass, but seed mass did not affect biomass response to offspring CO<sub>2</sub> treatments (no seed mass  $\times$  offspring CO<sub>2</sub> interaction was detected,  $F_{1,182} = 0.74$ ,  $P = 0.39$ ), and significant maternal CO<sub>2</sub>  $\times$  offspring CO<sub>2</sub> interactions were detected even when seed mass was included in the analyses as a covariate (Table 1). Mean family seed mass did not significantly influence any growth trait in *S. scoparium* or *P. pratensis* and also did not affect biomass responses to offspring eCO<sub>2</sub> treatments (seed mass  $\times$  offspring CO<sub>2</sub> interaction: *S. scoparium*  $F_{1,86} = 0.26$ ,  $P = 0.61$ ; *P. pratensis*  $F_{1,119} = 2.01$ ,  $P = 0.16$ ), although seed mass affected *P. pratensis* survival (Table 2). Results from analyses including germination date and seed mass as covariates were very similar to those from analyses that did not include these covariates. Thus,

**Table 2** Effects of maternal CO<sub>2</sub> and N treatments, offspring CO<sub>2</sub> treatment and all interactions on *L. perennis*, *P. pratensis*, and *S. scoparium* survival

	<i>L. perennis</i>		<i>P. pratensis</i>		<i>S. scoparium</i>	
	$\chi^2$	<i>P</i> value	$\chi^2$	<i>P</i> value	$\chi^2$	<i>P</i> value
CO <sub>2</sub> Offspring (O)	0.01	0.90	10.04	0.0015**	0.75	0.39
CO <sub>2</sub> Maternal (M)	0.02	0.90	0.48	0.49	1.06	0.30
N <sub>Maternal</sub>	4.09	0.04*	2.46	0.12	0.20	0.66
CO <sub>2</sub> Off × CO <sub>2</sub> M	0.33	0.57	0.01	0.93	0.81	0.37
CO <sub>2</sub> Off × N <sub>M</sub>	0.00	0.96	0.48	0.49	8.98	0.0027**
CO <sub>2</sub> M × N <sub>M</sub>	0.75	0.39	0.14	0.71	2.30	0.13
CO <sub>2</sub> Off × CO <sub>2</sub> M × N <sub>M</sub>	2.62	0.11	0.10	0.75	4.92	0.0265*
Germination date	3.37	0.07	32.20	<0.0001****	0.01	0.93
Seed mass	0.00	0.99	62.76	<0.0001****	0.98	0.32

\*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$ , \*\*\*\*  $P < 0.0001$

Values shown are chi-square statistics and associated *P* values from a logistic ANOVA. The proportion of germinates surviving to the final harvest was the response variable

these results are consistent with differences in seed mass between maternal treatments not being responsible for the differences in offspring responses to CO<sub>2</sub>.


## Discussion

Increases in resource availability as a result of rising atmospheric CO<sub>2</sub> concentrations and increased N deposition are expected to stimulate primary productivity. This prediction has been supported by field studies in a variety of natural communities, although biomass increases often depend on community composition and the availability of other nutrients (Reich et al. 2001a; Niklaus and Körner 2004; Norby et al. 2005; Rasse et al. 2005; see also Körner 2003). Because many growth chamber, greenhouse, and even field experiments involve only a single generation of plants, most studies are not of long enough duration to include the contribution of transgenerational effects to the magnitude of treatment response. However, as we have demonstrated here, increased resource availability can cause maternal effects or evolutionary responses that could either increase or decrease the plastic growth responses observed in single generation experiments.

*Lupinus perennis* and *P. pratensis* plants grown in eCO<sub>2</sub> were larger than those grown in aCO<sub>2</sub>, but plants grown from seeds collected from maternal eCO<sub>2</sub> environments responded less to offspring eCO<sub>2</sub> treatments than those grown from seeds collected from maternal aCO<sub>2</sub> environments. Thus, the transgenerational effect reduced the plastic phenotypic growth response to eCO<sub>2</sub>. Similar trends were observed for *S. scoparium*. Huxman et al. (2001) also found that maternal environmental effects of eCO<sub>2</sub> reduced the positive effects of offspring eCO<sub>2</sub> treatments on the growth

of *Bromus madritensis* ssp. *rubens*. While we observed few statistically significant transgenerational effects of N-addition on plant growth or survival, we detected some evidence that plant genotypes collected from N-addition maternal treatments showed greater (more positive) growth responses to offspring eCO<sub>2</sub> treatments than genotypes collected from ambient N maternal treatments. For *P. pratensis* survival, however, the opposite pattern was observed, and genotypes from N-addition maternal treatments exhibited lower survival under offspring eCO<sub>2</sub> treatments, while genotypes from ambient N maternal treatments experienced increased survival in eCO<sub>2</sub> offspring treatments.


Transgenerational effects may influence long-term responses to global environmental changes. Based on within-generation plastic growth responses to eCO<sub>2</sub>, eCO<sub>2</sub> is predicted to increase primary productivity (i.e., the “eCO<sub>2</sub> fertilization effect”). This increased productivity is expected to increase the effectiveness of the terrestrial CO<sub>2</sub> sink, thereby slowing the rate of increase in atmospheric CO<sub>2</sub> concentration (Hungate et al. 1997; Cramer et al. 2001). The transgenerational effects that we observed, however, suggest that the magnitude of the growth response to eCO<sub>2</sub> will decrease over time. Thus, estimates of biomass increases based on short-term plastic growth responses may overestimate the magnitude of the CO<sub>2</sub> fertilization effect by as much as 50%. If N-deposition continues to increase along with atmospheric CO<sub>2</sub> concentrations, however, the positive transgenerational effects of N-addition on plant growth response to offspring eCO<sub>2</sub> treatments may ameliorate some of the decrease in CO<sub>2</sub> responsiveness—although the magnitude and direction of transgenerational effects differed across traits (growth vs. survival), and some of the positive transgenerational effects of N-addition on growth responses to eCO<sub>2</sub> may be offset by the negative



**Fig. 2** Effects of offspring CO<sub>2</sub> treatment on survival of *L. perennis* (a), *P. pratensis* (b), or *S. scoparium* (c) populations collected from maternal ambient CO<sub>2</sub> (aCO<sub>2</sub>; open symbols) vs. elevated CO<sub>2</sub> (eCO<sub>2</sub>; filled symbols) and N-addition (squares) vs. ambient N (circles) treatments. Values shown are raw means ± 1SE. Symbols are offset so that non-overlapping error bars can be easily observed

transgenerational effects of N-addition on survival in eCO<sub>2</sub> offspring environments.

Because all seeds were germinated in a common greenhouse environment, the effects of offspring CO<sub>2</sub> environment on germination and early growth may have been missed. It is our belief, however, that germinating all seeds in a common, relatively benign environment likely had minimal impacts on the observed maternal environment × offspring environment interactions, especially since the magnitude of the maternal effects increased over time (data not shown). Additionally,



**Fig. 3** Effects of maternal CO<sub>2</sub> environment on the percentage biomass stimulation effect of offspring eCO<sub>2</sub> treatments for *L. perennis*, *S. scoparium*, or *P. pratensis*. For all species, genotypes from aCO<sub>2</sub> maternal environments (white bars) tended to respond more to offspring eCO<sub>2</sub> treatments than genotypes from eCO<sub>2</sub> maternal environments (black bars)

because we used field-collected seeds that were not propagated for one or more generations in a common environment, we were unable to differentiate between maternal environmental effects and evolutionary responses. Including seed mass and germination date in our analyses did not qualitatively alter the patterns that we observed, suggesting that the observed transgenerational effects were not primarily driven by differences in seed provisioning or germination timing. Seed nutrient concentrations were not measured, however, and are also potential sources of maternal environmental effects. Seeds produced from plants reared under eCO<sub>2</sub> and ambient N conditions might be expected to have lower nutrient (N) concentrations that could result in reduced responsiveness to eCO<sub>2</sub>. It is also plausible that genetic changes that have occurred over the five growing seasons of enhanced CO<sub>2</sub> and N also contributed to changes in eCO<sub>2</sub> responsiveness: plant populations growing in the long-term CO<sub>2</sub> and N treatments could have diverged genetically if N or CO<sub>2</sub> alter patterns of natural selection or the expression of genetic variation. While limited work has investigated how soil N influences predict evolutionary trajectories, a few examples suggest that eCO<sub>2</sub> can influence plant evolution, although the magnitude of these evolutionary effects is often weak (Bazzaz et al. 1995; Steinger et al. 2007). In contrast, a study by Lau et al. (2007) detected little evidence for direct evolutionary effects of eCO<sub>2</sub>.

In our experiment, species representing different functional groups seemed to differ in the magnitude of the transgenerational effect. There are at least three potential explanations for these differences among species. First, species differ in the magnitude of ecological response to eCO<sub>2</sub> (reviewed in Poorter and Navas 2003). In our experiment, we detected no evidence that eCO<sub>2</sub> stimulated the growth of *S. scoparium*, a C<sub>4</sub> plant. In contrast, both *L. perennis* and

*P. pratensis* responded positively to eCO<sub>2</sub>. If there is no plastic growth response to eCO<sub>2</sub>, then it seems less plausible that eCO<sub>2</sub> will alter resource allocation to seeds, thus minimizing the potential for maternal environmental effects. Second, species differed in the expression of family-level variation of growth traits and growth response to CO<sub>2</sub>. Maternal family explained significant amounts of variation in three of four measured *L. perennis* growth traits (Table 1); in contrast, significant family effects were detected in *P. pratensis* and *S. scoparium* only for culm heights (Table 1). These results suggest that while *L. perennis* may possess the genetic variation required for rapid evolutionary responses, genetic variation for most *P. pratensis* and *S. scoparium* growth traits may be lacking in these populations. Finally, interspecific differences in mating system and gene flow, life history, and generation time could all have influenced the likelihood of evolutionary change occurring over the relatively small temporal and spatial scales that these treatments have been applied.

Interestingly, Bezemer et al. (1998) document transgenerational effects of eCO<sub>2</sub> on the C3 grass *Poa annua* that are in the opposite direction of the transgenerational effects found in our study. While basic biological differences between *P. annua* and the species used in this study may contribute to these contrasting results, methodological differences, including differences in the duration of maternal treatments, could also play a role. In the Bezemer et al. (1998) study, maternal treatments were applied for a single generation, and seeds used in the offspring generation were collected from all plants in the population (not just the most fecund). Thus, the intensity of selection was relatively low, the potential for genetic change during the experiment was minimal, and the observed transgenerational effect was likely primarily a maternal environmental effect. In our study, genetic changes that may have occurred in response to five growing seasons of eCO<sub>2</sub> and N-addition treatments also could have contributed to the observed transgenerational effects and potentially could have counteracted any maternal environmental effects that were in the opposing direction.

In conclusion, our results suggest that transgenerational effects of maternal CO<sub>2</sub> environments may decrease plant growth responses to future eCO<sub>2</sub> conditions. Predictions of the long-term effects of eCO<sub>2</sub> on natural communities are typically based on biomass responses to short-term treatments that cannot assess transgenerational effects. Ignoring these multi-generation effects, however, could potentially bias predictions of future biomass and ecosystem responses to eCO<sub>2</sub> or other global changes.

**Acknowledgements** We thank A. Mueller for her assistance in field and R. Shaw, J. Dechaine, and two anonymous reviewers for providing helpful comments on an earlier draft of this manuscript. This project

was funded primarily by NSF IOB 0417094 to P. Tiffin, R. Shaw, and P. Reich and secondarily by NSF LTER (DEB 0080382) and Biocomplexity (0322057) programs, and a University of Minnesota Initiative on Renewable Energy and the Environment seed grant. This is KBS contribution #1462. The experiments described herein comply with current laws of the United States of America.

## References

- Aarssen LW, Burton SM (1990) Maternal effects at four levels in *Senecio vulgaris* (Asteraceae) grown on a soil nutrient gradient. *Am J Bot* 77:1231–1240
- Agrawal AA, Laforsch C, Tollrian R (1999) Transgenerational induction of defences in animals and plants. *Nature* 401:60–63
- Bazzaz FA (1990) The response of natural ecosystems to the rising global CO<sub>2</sub> levels. *Annu Rev Ecol Syst* 21:167–196
- Bazzaz FA, Jasienski M, Thomas SC, Wayne P (1995) Microevolutionary responses in experimental populations of plants to CO<sub>2</sub>-enriched environments: parallel results from two model systems. *Proc Natl Acad Sci USA* 92:8161–8165
- Bezemer TM, Thompson LJ, Jones TH (1998) *Poa annua* shows intergenerational differences in response to elevated CO<sub>2</sub>. *Glob Chang Biol* 4:687–691
- Collins S, Sultemeyer D, Bell G (2006) Changes in C uptake in populations of *Chlamydomonas reinhardtii* selected at high CO<sub>2</sub>. *Plant Cell Environ* 29:1812–1819
- Cramer W, Bondeau A, Woodward FI, Prentice IC, Betts RA, Brovkin V, Cox PM, Fisher V, Foley JA, Friend AD, Kucharik C, Lomas MR, Ramankutty N, Sitch S, Smith B, White A, Young-Molling C (2001) Global response of terrestrial ecosystem structure and function to CO<sub>2</sub> and climate change: results from six dynamic global vegetation models. *Glob Chang Biol* 7:357–373
- Durrant A (1958) Environmental conditioning of flax. *Nature* 181:928–929
- Ginzburg LR, Taneyhill DE (1994) Population cycles of forest lepidoptera: a maternal effect hypothesis. *J Anim Ecol* 63:79–92
- Hairston NG, Ellner SP, Geber MA, Yoshida T, Fox JA (2005) Rapid evolution and the convergence of ecological and evolutionary time. *Ecol Lett* 8:1114–1127
- Hungate BA, Holland EA, Jackson RB, Chapin FS, Mooney HA, Field CB (1997) The fate of carbon in grasslands under carbon dioxide enrichment. *Nature* 388:576–579
- Huxman TE, Hamerlynck EP, Jordan DN, Salsman KJ, Smith SD (1998) The effects of parental CO<sub>2</sub> environment on seed quality and subsequent seedling performance in *Bromus rubens*. *Oecologia* 114:202–208
- Huxman TE, Charlet TN, Grant C, Smith SD (2001) The effects of parental CO<sub>2</sub> and offspring nutrient environment on initial growth and photosynthesis in an annual grass. *Int J Plant Sci* 162:617–623
- Körner C (2003) Ecological impacts of atmospheric CO<sub>2</sub> enrichment on terrestrial ecosystems. *Philos Trans R Soc Lond B* 361:2023–2041
- Lau JA, Shaw RG, Reich PB, Shaw RG, Tiffin P (2007) Strong ecological but weak evolutionary effects of elevated CO<sub>2</sub> on a recombinant inbred population of *Arabidopsis thaliana*. *New Phytol* 175:351–362
- Lau JA, Strebom J, Stone LR, Reich PB, Tiffin P (2008) Direct and indirect effects of CO<sub>2</sub>, nitrogen, and community diversity on plant–enemy interactions. *Ecology* 89:226–236
- Niklaus PA, Körner C (2004) Synthesis of a six-year study of calcareous grassland responses to in situ CO<sub>2</sub> enrichment. *Ecol Monogr* 74:491–511

- Norby RJ, DeLucia EH, Gielen B, Calfapietra C, Giardina CP, King JS, Ledford J, McCarthy HR, Moore DJP, Ceulemans R, De Angelis P, Finzi AC, Karnosky DF, Kubiske ME, Lukac M, Pregitzer KS, Scarascia-Mugnozza GE, Schlesinger WH, Oren R (2005) Forest response to elevated CO<sub>2</sub> is conserved across a broad range of productivity. *Proc Natl Acad Sci USA* 102:18052–18056
- Platenkamp GAJ, Shaw RG (1993) Environmental and genetic maternal effects on seed characters in *Nemophila menziesii*. *Evolution* 47:540–555
- Poorter H, Navas M-L (2003) Plant growth and competition at elevated CO<sub>2</sub>: on winners, losers and functional groups. *New Phytol* 157:175–198
- Rasse DP, Peresta G, Drake BG (2005) Seventeen years of elevated CO<sub>2</sub> exposure in a Chesapeake Bay Wetland: sustained but contrasting responses of plant growth and CO<sub>2</sub> uptake. *Glob Chang Biol* 11:369–377
- Reale D, McAdam AG, Boutin S, Berteaux D (2003) Genetic and plastic responses of a northern mammal to climate change. *Proc R Soc Lond B* 270:591–596
- Reich PB, Knops J, Tilman D, Craine J, Ellsworth D, Tjoelker M, Lee T, Wedin D, Naeem S, Bahauddin D, Hendrey G, Jose S, Wragge K, Goth J, Bengston W (2001a) Plant diversity enhances ecosystem responses to elevated CO<sub>2</sub> and nitrogen deposition. *Nature* 410:809–812
- Reich PB, Tilman D, Craine J, Ellsworth D, Tjoelker MG, Knops J, Wedin D, Naeem S, Bahauddin D, Goth J, Bengston W, Lee TD (2001b) Do species and functional groups differ in acquisition and use of C, N and water under varying atmospheric CO<sub>2</sub> and N availability regimes? A field test with 16 grassland species. *New Phytol* 150:435–448
- Reich PB, Hobbie SE, Lee T, Ellsworth DS, West JB, Tilman D, Knops JMH, Naeem S, Trost J (2006) Nitrogen limitation constrains sustainability of ecosystem response to CO<sub>2</sub>. *Nature* 440:922–925
- Roach DA, Wulff RD (1987) Maternal effects in plants. *Annu Rev Ecol Syst* 18:209–235
- Rossiter MC (1996) Incidence and consequences of inherited environmental effects. *Annu Rev Ecol Syst* 27:451–476
- Shaw RG, Mitchell-Olds T (1993) ANOVA for unbalanced data: an overview. *Ecology* 74:1638–1645
- Snaydon RW (1970) Rapid population differentiation in a mosaic environment. I. The response of *Anthoxanthum odoratum* populations to soils. *Evolution* 24:257–269
- Snaydon RW, Davies MS (1972) Rapid population differentiation in a mosaic environment. II. Morphological variation in *Anthoxanthum odoratum*. *Evolution* 26:390–405
- Snaydon RW, Davies TM (1982) Rapid divergence of plant populations in response to recent changes in soil conditions. *Evolution* 36:289–297
- Steinger T, Stephan A, Schmid B (2007) Predicting adaptive evolution under elevated atmospheric CO<sub>2</sub> in the perennial grass *Bromus erectus*. *Glob Chang Biol* 13:1028–1039
- Trombulak SC (1991) Maternal influence on juvenile growth rates in Belding ground squirrel (*Spermophilus beldingi*). *Can J Zool* 69:2140–2145
- Wieneke S, Prati D, Brandl R, Stocklin J, Auge H (2004) Genetic variation in *Sanguisorba minor* after 6 years in situ selection under elevated CO<sub>2</sub>. *Glob Chang Biol* 10:1389–1401
- Yoshida T, Jones LE, Ellner SP, Fussmann GF, Hairston NG Jr (2003) Rapid evolution drives ecological dynamics in a predator–prey system. *Nature* 424:303–306
- Yoshida T, Hairston NG Jr, Ellner SP (2004) Evolutionary tradeoff between defense against grazing and competitive ability in a simple unicellular alga, *Chlorella vulgaris*. *Proc R Soc Lond B* 271:1947–1953